



Research article

Effect of vehicular traffic on pollen size and viability of Apocynaceous plant species

Mandeep Kaur and Avinash Kaur Nagpal*

Department of Botanical and Environmental Sciences, Guru Nanak Dev University, Amritsar-143005, Punjab, India

*Corresponding Author: avnagpal@yahoo.co.in

[Accepted: 04 June 2017]

Abstract: Vehicular traffic adds up different kinds of contaminants like heavy metals, polyaromatic hydrocarbons, gaseous toxins etc. to the atmosphere. Pollen grains of different plants have been found to be sensitive to these pollutants, which show varied effects on their viability, and size and hence can be used as bioindicators of air pollution. Keeping this in mind, the present work was designed to measure the pollen viability and pollen size of three common roadside plants, namely *Alstonia scholaris*, *Nerium oleander* and *Tabernaemontana divaricata* belonging to Apocynaceae family growing at seven sites of Amritsar city with varying levels of traffic stress. Pollen viability and pollen size were studied using 2, 3, 5-triphenyl tetrazolium chloride (TTC) staining method. The percentage of pollen viability of *A. scholaris* ranged from 32.45% to 65.74%, for *N. oleander* it ranged from 30.3% to 45.11% and for *T. divaricata* it ranged from 14.31% to 49.62%. Pollen size ranged from 22.86 to 23.94 μm for *A. scholaris*, 29.58 to 32.17 μm for *N. oleander* and 29.83-32.43 μm for *T. divaricata*. Pollen viability was found to be inversely proportional to a load of traffic at a given site whereas no remarkable variation was found in pollen size. For *A. scholaris*, a significant negative correlation ($r = -0.762$, $p \leq 0.05$) was observed between pollen viability and traffic load, while for other two species correlation between pollen viability and traffic load was not statistically significant at $p \leq 0.05$.

Keywords: Air pollutants - Indicator - Traffic stress - Pollen viability - Pollen size - TTC.

[Cite as: Kaur M & Nagpal AK (2017) Effect of vehicular traffic on pollen size and viability of Apocynaceous plant species. *Tropical Plant Research* 4(2): 235–241]

INTRODUCTION

Vehicular exhaust adds up huge amounts of soot particles, smoke, poisonous gases (SO_2 , NO_2 , CO_2 , VOCs etc.), heavy metals and organic molecules on the roads all over the world. All these air pollutants are known to produce adverse effects on the health of plants, animals and humans (Rezaei *et al.* 2010, Atkinson *et al.* 2012). Environmental pollution may also play a significant role in increasing pollen allergenicity and hence asthma problems in humans residing at various polluted areas (Majd *et al.* 2004). Plants growing outdoors are one of the main and primary targets to all types of air pollutants (Shannigrahi *et al.* 2004). Plants can act as great candidates for reduction of air pollution at a particular place by absorption, deposition, accumulation and detoxification of harmful pollutants (Prajapati & Tripathi 2008).

Many studies have shown effects of automobile emissions on leaves, stomatal apertures, pollen viability and growth patterns of plants (Iannotti *et al.* 2000, Verma *et al.* 2006). One of the major consequences of air pollution is its effect on pollen grain size, shape, exine sculpturing, viability etc. (Iannotti *et al.* 2000, Duro *et al.* 2013, Kaur *et al.* 2016). The mature pollen grains which are released from the flower buds are dry and while absorbing humidity from the ambient air, they also absorb some contaminants which influence their viability hence, plant's reproductive system (Kaur *et al.* 2016). Due to lipophilic nature of pollen and its exine surface stiffness, gaseous/particulate and organic/non-organic pollutants generally get accumulated inside it and resulting in a decrease in their size and different morphological deformities (Bist *et al.* 2004). Pollen, being sensitive indicators of air pollution, can provide interesting and useful information about the adverse effects of air pollution on living systems (Gottardini *et al.* 2004, Noori *et al.* 2009, Yousefi *et al.* 2011, Kaur *et al.* 2016).

Reproductive success of any plant is mainly dependent on the quantity and quality of pollen grains produced. Contaminants like pesticides, heavy metals, fluorides, etc. can accumulate on the surface of pollen grains and make them active and sensitive bio-indicators of atmospheric pollution (Noori *et al.* 2009, Yousefi *et al.* 2011). The literature survey has revealed enormous reports on the impact of air pollution on pollen grains structure and exine properties but very less work is accessible on the use of pollen grains as bioindicators of air pollution.

For assessment of pollen viability and size, different methods such as *in vitro* germination tests and staining methods are being used. TTC staining method is one of the widely and fast used methods based on dehydrogenase enzymatic reaction in mitochondria of living cells (Firmage & Dafni 2001, Soares *et al.* 2013).

Amritsar, a fast growing city of Punjab (India), is witnessing an exponentially increasing number of vehicles on the roads during the last few decades. Myllyvirta & Dahiya (2015) in their recent report published by WHO reported Amritsar to be 13th out of 20 most polluted Indian cities and it was found that vehicular exhaust adds an annual average of 92 $\mu\text{g}\cdot\text{m}^{-3}$ of PM_{2.5} (Particulate matter) into the atmosphere. Since roadside trees/shrubs are recipients of air pollution; their pollen studies can give an estimate of the biological effects of air pollution (Gottardini *et al.* 2004). Hence, the aim of the present study was to estimate the effects of traffic stress on pollen size and viability of three species of ornamental plants belonging to family Apocynaceae has widely grown on the roadsides of Amritsar.

MATERIALS AND METHODS

Area of study




Figure 1. Map showing location of study sites: **1**, Botanical Garden, Guru Nanak Dev University (BGG); **2**, Khalsa College Campus (KCC); **3**, Guru Nanak Dev University Whole Campus (GNC); **4**, Kabir Park Colony (KPC); **5**, Lahori Gate Road (LGR); **6**, Grand Trunk Road (GTR); **7**, Ram Bagh Chowk (RBC). [Source: <http://www.indmaps.com/state-map/punjab/>; <https://www.google.co.in/maps/@31.6336711,74.8000795,13z?hl=en>]

Table 1. Pollen sampling sites under study.

S. No.	Sampling site with code	Description of site
1	Botanical Garden, Guru Nanak Dev University (BGG)	It covers whole area of Botanical Garden of Guru Nanak Dev University, Amritsar.
2	Khalsa College Campus (KCC)	It covers whole campus of Khalsa College
3	Guru Nanak Dev University Whole Campus (GNC)	It covers whole campus of Guru Nanak Dev University excluding Botanical Garden area
4	Kabir Park Colony (KPC)	Residential colony covering road streets and park areas
5	Lahori Gate Road (LGR)	Road stretch from Beri gate to Lahori Gate
6	Grand Trunk Road (GTR)	Road stretch from Chheharta to Amritsar Railway Station
7	Ram Bagh Chowk (RBC)	It covers area around Ram Bagh Chowk

Amritsar, one of the industrialized and rapidly growing cities, is situated at N 31.6340°, E 74.8723° of the Northwestern state (Punjab) of India. The city lies at an average elevation of 234 m above sea level and covers almost 5075 Km² area. Amritsar has a semiarid climate, with four seasons *i.e.* winter season (December to March) with minimum temperature of 0°C and maximum of about 15°C; summer season or pre-monsoon (April to June) where temperature can go up to 42°C; monsoon season from July to September while post-monsoon season is from October to November. Annual rainfall is about 681 millimetres. Its population is about 1,132,761 (Census of India 2011). In the present study, seven roadside sites of Amritsar city were selected (Fig. 1; Table 1).

Plants under study

Three apocynaceous plant species namely, *Alstonia scholaris* (L.) R. Br., *Nerium oleander* L. and *Tabernaemontana divaricata* (L.) R.Br. ex Roem. & Schult. growing along the roadsides of Amritsar city, Punjab, India were considered. Brief description of plants under study is presented in table 2. The reason for selection of these plants was their common occurrence at all the studied sites.

Table 2. Description of different plants under study.

Plant Name	General Features observed
Botanical name- <i>Alstonia scholaris</i> (L.) R. Br. Common name- Chattiyar/Indian devil tree Family- Apocynaceae	<ul style="list-style-type: none"> • Grows up to 100-130 ft, flowers bloom in the month of October, leaves are glossy and occur in whorls of 3-10; petioles are 1–3 cm. • In Ayurveda, it is an astringent herb for treating skin disorders, malarial fever, urticaria, chronic dysentery, diarrhea, snake bite and milk used to treat beriberi disease.
Botanical name- <i>Nerium oleander</i> L. Common name- Kaner/ Oleander Family- Apocynaceae	<ul style="list-style-type: none"> • Grows up to 10-13 ft in height, leaves are 10 to 22 cm long. The plant produces terminal flower heads, usually pink or white. Each flower is about 5 cm in diameter and five- petalled. • The whole plant, including the milky white sap, is toxic. The one of the most studied toxin is oleandrin.
Botanical name- <i>Tabernaemontana divaricata</i> (L.) R.Br. ex Roem. & Schult. Common name- Chandni/Crape Jasmine Family- Apocynaceae	<ul style="list-style-type: none"> • Grows to a height of 5 to 6 ft, leaves are 6 inches in length and 2 inches in width, flowers are pinwheel shaped. • Root decoction is used in treatment of ascariasis, scabies, internal poisoning, kidney stones and also used as antipyretic while stems are used to treat dental caries.

Collection of buds

Sampling was carried out once in a year, 2014. Flower buds of *N. oleander* and *T. divaricata* were collected during the months of May–July while for *A. scholaris* sampling for flower buds was done during October–November. Fresh and young flower buds were collected during early morning hours (5.30–8.30 am) from each species growing at each of the seven sites of Amritsar city. At least 10 flower buds of suitable size were collected from each plant. For each site of the study, buds from three plants of each species were collected. Experiments were carried out in triplicates.

Traffic load

The traffic load was calculated by counting the number of vehicles including heavy vehicles, four wheelers, three wheelers and 2 wheelers moving at each site for 3 h at different intervals of time (8–9 am; 12–1 pm; 4–5 pm) with the help of a hand-held counter.

Pollen viability

Pollen viability was estimated by employing 2, 3, 5-triphenyl tetrazolium chloride (TTC) staining method according to the protocol of Iannotti *et al.* (2000) with slight modifications. Freshly prepared TTC stain kept in dark bottles was used for making slides every time. TTC stain was prepared by taking 1% solution of TTC in 60% sucrose solution. A minimum of 10 anthers from at least 5–8 flower buds of suitable size of each plant were excised with a sharp needle and placed on a slide. Further, a drop of TTC stain was added to squash anthers on the glass slide. A cover slip was placed on the squashed material and air bubbles were removed. The slides were sealed with dibutylphthalate xylene (DPX) and incubated for 2 hrs in daylight. Pollen grains slides were prepared in the field or in laboratory immediately after sampling. The stained slides were then preserved in air tight box. After one day, slides were observed under a light microscope at 400X magnification at room temperature for viable and non-viable pollen grains. Pollen viability was calculated by counting viable and non-viable pollen grains. The deep pink or red stained pollen grains with round shape were counted as viable while white or yellowish pollen grains with shriveled structure were counted as non-viable. A minimum of 1000 pollen grains were studied for each plant. The experiment was carried out in triplicates.

Calculations

Pollen viability was calculated as,

$$\text{Pollen viability (\%)} = \frac{\text{Number of viable pollen grains}}{\text{Total number of counted pollen grains}} \times 100$$

Pollen size

Light microscope fitted with an ocular micrometer was used to study the size of pollen grains under the magnification of 400X. Ocular micrometer was calibrated by using stage micrometer. At least 10 pollen grains were observed for each plant species on average and the studies were carried out in triplicates.

Statistical analysis

Data was analysed by using the self-coded software on Microsoft Excel 2007 for one-way ANOVA test to compare differences among the studied variables. Tukey's Honestly Significant Difference test (Tukey's HSD) was also performed for multiple comparisons between the means. Correlations between the parameters were also evaluated by calculating the Pearson's correlation coefficient (r).

RESULTS

In the present study, an attempt was made to assess the effect of traffic load on pollen of three plant species growing at seven sites of Amritsar city under different levels of traffic. Results for traffic load, pollen viability and pollen size are presented in tables 3. Table 3 depicted varying levels of traffic for each site under study. The highest number of vehicles was observed at the site RBC whereas the lowest number was observed at site KCC. Though vehicles are not allowed inside Botanical Garden of Guru Nanak Dev University (BGG); however, it is adjacent to GT Road, NH 1 (Table 3).

The percentage of pollen viability for *Alstonia scholaris* ranged from 32.45% (GTR) to 65.74% (GNC), for *Nerium oleander* 30.3% (LGR) to 45.11% (KCC) and for *Tabernaemontana divaricata* 14.31% (RBC) to 49.62% (KPC). Pollen viability was found to be inversely proportional to the number of vehicles at a given site. Figure 2 shows viable and non-viable pollen grains of *A. scholaris*, *N. oleander* and *T. divaricata* plant species. Pollen size of *A. scholaris* was observed to range from 22.86 μm (GNC) to 23.94 μm (GTR), for *N. oleander* 29.58 μm (GTR) to 32.17 μm (RBC) and for *T. divaricata* it ranged from 29.83 μm (RBC) to 32.43 μm (KPC).




Figure 2. Viable and non-viable pollen grains of the plant species studied.

One way ANOVA test was used to compare the differences between the parameters observed for different plant species growing at different sites of samples collection. ANOVA for pollen viability showed the statistical significant difference with respect to their sites of samples collection whereas no statistically significant variation was observed in pollen size with respect to different sites studied (Table 3). Tukey's HSD test was also used to check the differences in pollen viability and pollen size at different sites studied.

Pearson correlation coefficient (r) was calculated to find out the correlation of pollen viability with traffic load and pollen size with traffic load at $p \leq 0.05$ level of significance (Table 3). In the case of *A. scholaris*, a significant negative correlation between pollen viability and traffic load at different sites was observed ($r = -$

0.762, $p \leq 0.05$), while other two species did not show any significant correlation. Pollen size did not show any significant correlation with traffic density for all studied species.

DISCUSSION

Automobile generated air pollutants and different environmental stresses have played a major role in influencing the pollen viability and pollen size of different plant species (Gottardini *et al.* 2004, Higashitani 2013, Rezanejad 2013, Paupière *et al.* 2014).

The viability of pollen grains of plants under study was determined by TTC staining method which works on the colouring of tissue. TTC gets reduced by the dehydrogenase enzyme of live tissues and results information of a carmine red colour compound known as formazan (Beyhan & Serdar 2008). This method is relatively simple and gives rapid results (Soares *et al.* 2013). This technique has been commonly used for studying the effects of automobile generated airborne pollutants including heavy metals on viability/fertility of pollen grains of several plant species (Duro *et al.* 2013). The results obtained by this method are well comparable to those obtained by *in vitro* germination tests (Huang *et al.* 2004). Many studies have suggested the use of TTC as a reliable method for assessment of pollen viability (Iannotti *et al.* 2000, Gottardini *et al.* 2004, Huang *et al.* 2004, Abdelgadir *et al.* 2012, Kaur *et al.* 2016). Staining techniques used for estimation of pollen viability and size of pollen grains depend on the type of plant species. These stains/dyes can easily point out differentiation among viable and non-viable pollen grains (Kolodziejek & Gabara 2008, Sharafi 2011).

In the present study, as a result of vehicular emissions, remarkable effects on pollen viability of three plant species under study were observed. Viable pollen grains were found to be darkly stained and round in shape while non-viable pollen grains were found to be transparent and have shriveled structure. Pollen viability of plant species studied was found to be negatively correlated with the traffic load (no. of vehicles) of different sites where these plants were growing. Effects of air pollutants on pollen viability and size have been well documented for different plant species (Chauhan & Singh 1996, Malayeri *et al.* 2012, Greguskova & Micieta 2013). Dickinson (2000) and Rezanejad (2009) observed alterations in proteins involved in pollen-stigma interactions as a consequence of automobiles generated air pollution.

Pollen size of *Alstonia scholaris*, *Nerium oleander* and *Tabernaemontana divaricata* at all sites was found to be unaffected irrespective of high vehicle number. One way ANOVA test showed no significant differences between pollen grains size from different sites of sample collection whereas pollen viability of *A. scholaris* ($p \leq 0.001$), *T. divaricata* ($p \leq 0.001$) and *N. oleander* ($p \leq 0.05$) showed statistically significant differences with respect to their sites of sample collection. Our results are in conformity with some of the earlier studies on different plant species like *Daucus carota* L., *Convolvulus sepium* L. and *Dactylis glomerata* L. (Iannotti *et al.* 2000); *Cassia fistula* L., *Cannabis sativa* L. and *Thevetia peruviana* (Pers.) K. Schum. (Kaur *et al.* 2016).

Pearson's Correlation Coefficient analysis suggests that among the three species studied, *Alstonia scholaris* ($r = -0.762$, at $p \leq 0.05$) was found to be most sensitive to air pollution followed by *Tabernaemontana divaricata* ($r = -0.645$); *Nerium oleander* ($r = -0.529$) was relatively tolerant to traffic stress (Table 3). Hence, *Alstonia scholaris* and *Tabernaemontana divaricata* can be used as better bio-indicators of vehicular air pollution whereas *N. oleander* can be the plant of choice for polluted sites. Earlier also it was reported that pollen viability can be used as reliable bio-indicator of atmospheric pollution (Iannotti *et al.* 2000, Greguskova & Micieta 2013). Malayeri *et al.* (2012) reported the use of pollen viability and morphology of six legumes *viz.*, *Cercis siliquastrum* L., *Medicago sativa* L., *Robinia pseudoacacia* L., *Melilotus officinalis* (L.) lam, *Trifolium repens* L., and *Sophora alba* L. for bio-monitoring of fluoride generated pollution.

CONCLUSION

The present study showed that pollen viability relatively decreases with increase in vehicle number while pollen size did not show any significant variation. On the basis of pollen viability data, *N. oleander* was found to be tolerant to air pollution while *A. scholaris* was found to be highly sensitive followed by *Tabernaemontana divaricata*, hence, the latter two can act as better indicators of air pollution with respect to traffic stress. Present work provides information for three species only, which is insufficient to draw a specific conclusion. The effect on pollen viability and pollen size depends on the type of plant species and level of traffic stress. It can be concluded that pollen grains can give important information about the biological impact of air pollutants and can be used as good candidates as bio-indicators of the atmospheric pollution. This kind of work may be helpful in exploration of tolerance or sensitivity level of different roadside plant species against air pollution.

ACKNOWLEDGEMENTS

Thanks are due to University Grants Commission (UGC) for financial assistance under CPEPA, UPE and DRS programmes. Mandeep Kaur is recipient of UGC-BSR fellowship. Authors also thank Prof. A. K. Thukral, Department of Botanical & Environmental Sciences, GNDU for designing self coded software in MS Excel 2007 for statistical analysis of the data.

REFERENCES

- Abdelgadir HA, Johnson SD & VanStaden J (2012) Pollen viability, pollen germination and pollen tube growth in the biofuel seed crop *Jatropha curcas* (Euphorbiaceae). *South African Journal of Botany* 79: 132–139.
- Atkinson RW, Yu D, Armstrong BG, Pattenden S, Wilkinson P, Doherty RM, Heal MR & Anderson HR (2012) Concentration–Response function for Ozone and daily mortality: Results from five urban and five rural U.K. populations. *Environmental Health Perspectives* 120 (10): 1411–1417.
- Beyhan N & Serdar U (2008) Assessment of pollen viability and germinability in some European chestnut genotypes (*Castanea sativa* L.). *Horticultural Science (Prague)* 35(4): 171–178.
- Bist A, Pandit T, Bhatnagar AK & Singh AB (2004) Variability in protein content of pollen of Castor bean (*Ricinus communis*) before and after exposure to the air pollutants SO₂ and NO₂. *Grana* 43: 94–100.
- Chauhan SVS & Singh J (1996) Deterioration of pollen fertility in response to air pollution: A review. In: Agrawal SK, Tiwari S & Dubey PS (eds) *Biodiversity and Environment*. APH Publ. Co, New Delhi, pp. 197–205.
- Dickinson HG (2000) Pollen stigma interactions: So near yet so. *Trends in Genetics* 16: 373–376.
- Duro A, Piccione V & Zampino D (2013) Air quality biomonitoring through pollen viability of Fabaceae. *Environmental Monitoring and Assessment* 185: 3803–3817.
- Firmage DH & Dafni A (2001) Field tests for pollen viability; a comparative approach. *Acta Horticulturae* 561: 87–94.
- Gottardini E, Cristofolini F, Paoletti E, Lazzeri P & Pepponi G (2004) Pollen viability for air pollution biomonitoring. *Journal of Atmospheric Chemistry* 49: 149–159.
- Greguskova E & Micieta K (2013) Phytoindication of the ecogenotoxic effects of vehicle emissions using pollen abortion test with native flora. *Polish Journal of Environmental Studies* 22 (4): 1069–1076.
- Higashitani A (2013) High temperature injury and auxin biosynthesis in microsporogenesis. *Frontier of Plant Sciences* 4: 1–4.
- Huang Z, Zhu J, Mu X & Lin J (2004) Pollen dispersion, pollen viability and pistil receptivity in *Leymus chinensis*. *Annals of Botany* 93: 295–301.
- Iannotti O, Mincigrucci G, Bricchi E & Frenguelli G (2000) Pollen viability as a bio- indicator of air quality. *Aerobiologia* 16: 361–365.
- Kaur M, Sharma A, Kaur R, Katnoria JK & Nagpal AK (2016) Palynological studies of some roadside plants under exposure to traffic stress. *Aerobiologia* 32 (2): 245–254.
- Kolodziejek J & Gabara B (2008) Palynological study of Polish taxa of *Potentilla* subsect. *Collinae* (Rosaceae). *Acta Botanica Croatica* 67 (2): 139–146.
- Majd A, Chehregani A, Moin M, Gholami M, Kohno S, Nabe T & Shariatzade MA (2004) The effect of air pollution on structures proteins and allergenicity of pollen grains. *Aerobiologia* 20: 111–118.
- Malayeri BE, Noori M & Jafari M (2012) Using the pollen viability and morphology for fluoride pollution biomonitoring. *Biology and Trace Element Research* 147: 315–319.
- Myllyvirta L & Dahiya S (2015) A status assessment of National Air Quality Index (NAQI) and pollution level assessment for Indian cities. A report by WHO. Published by Greenpeace, India, pp. 1–19.
- Noori M, Malayeri BE & Jafari M (2009) Determination of fluoride and its effects on flavonoids in some legumes. *Toxicology and Environmental Chemistry* 91 (3): 409–418.
- Paupière MJ, van Heusden AW & Bovy AG (2014) The metabolic basis of pollen thermo-tolerance: *Perspectives for Breeding. Metabolites* 4: 889–920.
- Prajapati SK & Tripathi BD (2008) Seasonal variation of leaf dust accumulation and pigment content in plant species exposed to urban particulates pollution. *Journal of Environmental Quality* 37: 865–870.
- Census of India (2011) Cities having population 1 lakh and above. In: Provisional Population Totals. Office of the Registrar General & Census Commissioner, India. Available from: <http://www.censusindia.gov.in/pca/SearchDetails.aspx?Id=42494> (accessed: 01 Oct. 2016).

- Rezaei M, Arzani A & Sayed-Tabatabaei BE (2010) Meiotic behavior of tetraploid wheats (*Triticum turgidum* L.) and their synthetic hexaploid wheat derivatives influenced by meiotic restitution and heat stress. *Journal of Genetics* 89: 401–407.
- Rezanejad F (2009) Air pollution effects on structure, proteins and flavonoids in pollen grains of *Thuja orientalis* L. (Cupressaceae). *Grana* 48 (3): 205–213.
- Rezanejad F (2013) The response of anther and pollen development, pollen cellular material release and pollen proteins to air pollution in *Petunia* hybrid Juss. (Solanaceae). *Iranian Journal of Science and Technology, Transaction A: Science* 1: 63–68.
- Shannigrahi AS, Fukushima T & Sharma RC (2004) Anticipated air pollution tolerance of some plant species considered for green belt development in and around an industrial/urban area in India: an overview. *International Journal of Environmental Studies* 61 (2): 125–137.
- Sharafi Y (2011) An investigation on the pollen germination and tube growth in some *Prunus persica* genotypes and cultivars. *African Journal of Microbiology Research* 5 (14): 2003–2007.
- Soares TL, de Jesus ON, de Souza EH, dos Santos-Serejo JA & de Oliveira EJ (2013) Morphology and viability of pollen grains from passion fruit species (*Passiflora* spp.). *Acta Botanica Brasiliica* 27(4): 779–787.
- Verma RB, Mahmooduzzafar, Siddiqi TO & Iqbal M (2006) Foliar response of *Ipomea pestigridis* L. to coal smoke pollution. *Turkish Journal of Botany* 30: 423–417.
- Yousefi N, Chehregani A, Malayeri BE, Lorestani B & Cheraghi M (2011) Investigating the effect of heavy metals on developmental stages of anther and pollen in *Chenopodium botrys* L. (Chenopodiaceae). *Biology and Trace Element Research* 140: 368–376.