



Research article

## Assessment of diversity, population structure and regeneration status of tree species in Hollongapar Gibbon Wildlife Sanctuary, Assam, Northeast India

Moumita Sarkar and Ashalata Devi\*

Department of Environmental Science, Tezpur University, Tezpur, Sonitpur, Assam, India

\*Corresponding Author: [ashalatadevi12@gmail.com](mailto:ashalatadevi12@gmail.com)

[Accepted: 20 August 2014]

**Abstract:** The present study was carried out for quantitative analysis of diversity, population structure and regeneration status of tree species in tropical semi-evergreen forest of Hollongapar Gibbon Wildlife Sanctuary, Assam, northeast India. The study was conducted during 2010–2011, by laying 100 quadrats (10×10 m) following random plot sampling method. A total of 75 tree species ( $\geq 30$  cm gbh), belonging to 60 genera and 40 families were recorded from the study area. Individuals were categorized into three groups, seedling, sapling and adult based on girth classes and the status of natural regeneration of species was determined based on their population size. Highest density (7756 individuals.ha<sup>-1</sup>) and species richness (73) were recorded in 0–30 cm girth class, while highest basal area (9.62 m<sup>2</sup>.ha<sup>-1</sup>) was observed in 120–150 cm girth class. Majority of tree species (36%) exhibited ‘fair regeneration’ condition followed by ‘good regeneration’ status (24%). The overall population structure of tree species shows a reverse J-shaped population curve and ‘good’ regeneration status which reveals that the future communities may be sustained. The study gives an understanding of the diversity, pattern of population and regeneration of the tree species of the sanctuary which may help in forest management and conservation of the species.

**Keywords:** Conservation - Forest - Population structure - Protected areas - Quantitative characters.

[Cite as: Sarkar M & Devi A (2014) Assessment of diversity, population structure and regeneration status of tree species in Hollongapar Gibbon Wildlife Sanctuary, Assam, Northeast India. *Tropical Plant Research* 1(2): 26–36]

### INTRODUCTION

The structure and function of forest ecosystem is determined by the plant component more than any other living component of the system (Richards 1996). The plant diversity at any site is influenced by species distribution and abundance patterns (Palit & Chanda 2012) and the richness of plant species is controlled by a variety of biotic and abiotic parameters (Rannie 1986, Huston 1994). Topography, soil, climate and geographical location of a region influence the vegetation diversity of the forest ecosystem (Ram *et al.* 2004). It was found that, plant diversity inventories in tropical forests have mostly been concentrated on tree species than other life-forms (Mani & Parthasarathy 2006). The nature of forest communities largely depends on the ecological characteristics in sites, species diversity and regeneration status of species (Khumbongmayum *et al.* 2006). Species diversity is one of the most important indices used for evaluating the stability and sustainability of forest communities. Information on the species composition of a forest is essential for its wise management in terms of economic value, regeneration potential (Wyatt-Smith 1987) and ultimately may be leading to conservation of biological diversity (Verma *et al.* 1999).

Population structure is expressed in terms of number of individuals present in each of the definite girth class distribution of tree species. Saxena & Singh (1984) reported regeneration behaviour of tree species in a forest can be revealed from the population structure. A successful regeneration is indicated by presence of sufficient number of seedlings, saplings and young trees in a given population (Pokhriyal *et al.* 2010) and the number of seedling of any species can be considered as the regeneration potential of that species (Negi & Nautiyal 2005). Natural regeneration is a central component for tropical forest ecosystem dynamics (Getachew *et al.* 2010) and is essential for preservation and maintenance of biodiversity (Rahman *et al.* 2011). It is important to understand the growth status of a species in the ecosystem and is one of the key parameter to determine ecosystem stability (Kadavul & Parthasarathy 2001, Deb & Sundriyal 2011). Several types of disturbances like logging, landslides,

gap formation, litterfall, herbivory, etc. can affect the potential regenerative status of species composing the forest stand spatially and temporally (Guariguata 1990, Welden *et al.* 1991, Barik *et al.* 1996a, Boerner & Brinkman 1996, Liang & Seagle 2002, Ganesan & Davidar 2003, Khumbongmayum *et al.* 2005, Ceccon *et al.* 2006, Khumbongmayum *et al.* 2006, Ward *et al.* 2006, Guarino & Scariot 2012).

The northeast India is a storehouse of rich biodiversity which includes variety of plant and animal species and it is considered as one of the richest biodiversity centres of the Indian continent (Tynsong & Tiwari 2010) with rich species density and diversity (Nath *et al.* 2005). Assam, a state in northeast India, has total recorded forest area of 28,748 km<sup>2</sup>. which covers 32% of the total geographical area of the state and harbours 3017 species of flowering plants (Patiri & Borah 2007). Most of the population and regeneration studies in Northeast India were reported from the states of Arunachal Pradesh, Meghalaya, Manipur, Mizoram and Tripura, but a few studies from Assam (Borah & Garkoti 2011, Nandy & Das 2013, Dutta & Devi 2013, Saikia & Khan 2013). Therefore, the present study was undertaken to analyse the tree species diversity, population dynamics and to assess the regeneration pattern of tree species of this tropical semi-evergreen forest of Assam, Northeast India. The findings of the study will definitely add records on quantitative data on tree species diversity of forest of Assam in particular and tropical forest in general.

## MATERIAL AND METHODS

### Study area

Hollongapar Gibbon Wildlife Sanctuary (26°40'–26°45' N and 94°20'–94°25' E) is situated in Mariani range of Jorhat district in upper Assam at an altitudinal range of 100–120 m above msl. It has got the status of Reserve forest in 1881 and then in 1997, the Hollongapar Reserve Forest was upgraded to Gibbon Wildlife Sanctuary (Fig. 1). Again in the year 2004, its name was changed to Hollongapar Gibbon Wildlife Sanctuary (abbreviated as HGWLS hereafter) but it is still popularly known as Gibbon Wildlife Sanctuary. The area is situated amidst tea gardens and villages, crisscrossed with numerous rain fed streams (*nallahs*). It covers an area of 19.49 km<sup>2</sup> and the sanctuary has been divided into five compartments. The area receives an average annual rainfall of 249 cm. The soil is sandy clay loam in texture, slightly acidic in nature having pH 5.1 and soil organic carbon content records 2.03%. The weather in the area may be classified as subtropical hot, wet monsoon periods (May–August) and cool dry winter (September to April). Winter rains are also not uncommon and the average temperature varies from 5°C (min) – 38°C (max).




Figure 1. Map of the study site.


As per Champion & Seth (1968), classification scheme, the forest type of HGWLS is “Assam Plains Alluvial Semi Evergreen Forests (1/2/2B/C)”. The sanctuary is a suitable habitat for large number of mammalian, birds and invertebrate species. It harbours seven species of primates viz. Western Hoolock gibbon, Slow loris, Capped langur, Rhesus, Pigtailed, Stump-tailed and Assamese macaque. This Sanctuary has the rare distinction of holding one of the highest densities of gibbon populations in Assam. Extensive studies on primates have been carried out in this sanctuary by several workers, but there is a lack of baseline information and detailed study on quantitative characteristics of diversity, population structure and natural regeneration status of tree species of HGWLS. Therefore, the study has been carried out and the recorded data on quantitative characters of tree species may be helpful to formulate conservation strategy for plant species in particular and for the proper conservation of animal species inhabiting in the sanctuary in general.

### Methods

The study for the assessment of tree diversity, population structure and quantitative characteristic features was conducted during 2010–2011 using random sampling method. Extensive field survey in all the five compartments was carried out in the sanctuary during phytosociological study period. Quadrat method was followed to record the tree species diversity and other quantitative parameters. For the study, 100 quadrats of 10×10 m were laid down randomly in the study site, covering an area of 1 ha. All the species and individuals encountered in each quadrat were counted and the girth was measured. Individuals having  $\geq 30$  cm girth (gbh) were considered as adult, saplings with  $\geq 10$  cm to  $< 30$  cm girth and seedlings with  $< 10$  cm girth. The number of individuals and girth of each individual species encountered in each quadrat were used for further quantitative analysis. Population structure of the species was analyzed across thirteen girth classes *i.e.* 0–30, 30–60, 60–90, 90–120, 120–150, 150–180, 180–210, 210–240, 240–270, 270–300, 300–330, 330–360 and  $>360$  cm. The status of regeneration of species was determined based on population size of seedlings, saplings and adults as (modified from Khan *et al.* 1987, Shankar 2001, Khumbongmayum *et al.* 2006): (a) ‘good’, if seedlings  $>$  or  $<$  saplings  $>$  adults; (b) ‘fair’, if seedlings  $>$  or  $\leq$  saplings  $\leq$  adults; (c) ‘poor’, if a species survives only in sapling stage, but no seedlings (though saplings may be  $<$ ,  $>$  or  $=$  adults); (d) ‘none’, if it is absent both in sapling and seedlings stages, but found only in adults and (e) ‘new’, if a species has no adults, but only saplings and/or seedlings. Community quantitative parameters such as frequency, density, abundance, basal area (BA), relative frequency, relative density, relative dominance and Importance Value Index (IVI) were calculated (Cottam & Curtis 1956). The Shannon-Wiener index ( $H'$ ) (Shannon & Weaver 1963), Simpson’s index ( $C_D$ ) (Simpson 1949) and Pielou’s evenness index ( $e$ ) (Pielou 1966) were also evaluated. Identification of plants was made by referring taxonomic literature (Kanjilal & Bor 2005) and by consulting plant specimen at Botanical Survey of India, Itanagar, Arunachal Pradesh. Herbarium specimen are deposited and preserved in the Department of Environmental Science, Tezpur University.

## RESULTS

### Tree diversity




**Figure 2.** Tree species: **A**, Dominance diversity (D-D) curve; **B**, Density, basal area and species richness in different girth classes.

A total of 75 tree species, belonging to 60 genera under 40 families were recorded from the study area. Moraceae was the dominant family having 8 species followed by Magnoliaceae with 5 species, Anacardiaceae,

Euphorbiaceae, Lauraceae and Meliaceae recorded 4 species each. Under Moraceae family, the genus *Ficus* contributed highest number of six species viz. *Ficus benghalensis*, *Ficus benjamina*, *Ficus fistulosa*, *Ficus lamponga*, *Ficus racemosa* and *Ficus religiosa*. The total basal area of the tree species was computed as 58.0 m<sup>2</sup>.ha<sup>-1</sup> with density of 750 individuals.ha<sup>-1</sup>. The dominant tree species was *Vatica lanceaefolia* Bl. (Dipterocarpaceae), a critically endangered species as given by International Union for Conservation of Nature red list of threatened species (IUCN 2014), which contributed 55.52 IVI value recording highest density (227 individuals.ha<sup>-1</sup>) with basal area of 6.525 m<sup>2</sup>.ha<sup>-1</sup>. Detailed quantitative data of each tree species, density (ha<sup>-1</sup>), basal area (m<sup>2</sup>.ha<sup>-1</sup>) and IVI values are given in table 1. The other important species based on IVI values were *Artocarpus chaplasha* (18.13), *Lagerstroemia speciosa* (16.30), *Magnolia hookeri* (16.07) and *Dipterocarpus retusus* (13.91) as shown in fig. 2A. The Shannon-Wiener index (H'), Simpson's index (C<sub>D</sub>) and Evenness index (e) for the tree species were calculated as 3.55, 0.05 and 0.82, respectively.

#### Population structure and regeneration status

Overall population structure of tree species depending on size-class distribution yielded reverse J-shaped curve in HGWLS (Fig. 2B). The highest percentage (91.22%) of tree individuals were recorded in 0–30 cm girth class and it gradually decreased with increasing girth class. Highest density (7756 individuals.ha<sup>-1</sup>) and species richness (73) were recorded in 0–30 cm girth class, while *Tetrameles nudiflora* showed lowest density (1 individual.ha<sup>-1</sup>) and species richness (1) in 330–360 cm girth class. Highest basal area (9.62 m<sup>2</sup>.ha<sup>-1</sup>) was observed in 120–150 cm girth class and lowest (0.97 m<sup>2</sup>.ha<sup>-1</sup>) in 330–360 cm girth class. Population structure of a few dominant tree species in HGWLS such as *Magnolia hookeri*, *Dipterocarpus retusus*, *Artocarpus chaplasha* and *Vatica lanceaefolia* showed reverse J-shaped population curve, whereas *Lagerstroemia speciosa* showed an interrupted population curve (Fig. 3).



**Figure 3.** Population structure of few dominant tree species recorded in the study site: **A**, *Magnolia hookeri*; **B**, *Dipterocarpus retusus*; **C**, *Artocarpus chaplasha*; **D**, *Vatica lanceaefolia*; **E**, *Lagerstroemia speciosa*.

In the present study, 24% tree species exhibited ‘good’ regeneration status, 36% showed ‘fair’ regeneration condition and 8% showed ‘poor’ regeneration status. A total of 17% tree species were ‘not regenerating’ at all and 15% tree species, which were available only in sapling or seedling stage, were considered as ‘new’ in HGWLS (Table 1). The ‘poor’ regenerating tree species were *Aglaia spectabilis*, *Cinnamomum glaucescens*, *Dillenia indica*, *Ficus racemosa*, *Hydnocarpus kurzii*, *Magnolia hodgsonii* and *Terminalia catappa*. Species which were found in ‘none’ or ‘not regenerating’ category were *Albizia lebbek*, *Alstonia scholaris*, *Cyathea gigantea*, *Duabanga grandiflora*, *Evodia meliaefolia*, *Macaranga denticulate*, *Morinda angustifolia*, *Neolamarckia cadamba*, *Palaquium obovatum*, *Pterospermum acerifolium*, *Quercus gemelliflora*, *Sterculia villosa*, *Terminalia myriocarpa*, *Vernonia arborea* and *Walsura robusta*. ‘New’ regeneration status included

*Actinodaphne angustifolia*, *Albizia lucidior*, *Bischofia javanica*, *Chrysophyllum roxburghii*, *Cinnamomum bejolghota*, *Gynocardia odorata*, *Horsfieldia kingii*, *Machilus gamblei*, *Meliosma pinnata*, *Phoebe goalparensis*, *Syzygium cumini*, *Terminalia bellirica* and *Toona ciliata*. The total density of seedlings (6754 individuals.ha<sup>-1</sup>) was recorded to be higher than the saplings (1002 individuals.ha<sup>-1</sup>) and adults (750 individuals.ha<sup>-1</sup>) in the study site, thus exhibiting overall ‘good’ regeneration condition. The dominant tree species, *Vatica lanceaefolia* had maximum seedling density (2876 individuals ha<sup>-1</sup>) with highest sapling and adult density of 412 and 227 individuals ha<sup>-1</sup>, respectively showing ‘good’ regeneration status. Among all tree species recorded in the study site, *Vatica lanceaefolia* is a critically endangered species which contributed 37% towards the overall regeneration status.

**Table 1.** Density (D=individuals.ha<sup>-1</sup>), basal area (BA=m<sup>2</sup>.ha<sup>-1</sup>), Importance Value Index (IVI) and regeneration status (RS) of adult tree species of Hologapar Gibbon Wildlife Sanctuary.

Scientific Name	Family	D	BA	IVI	RS
<i>Actinodaphne obovata</i> (Nees.) Bl.	Lauraceae	17	0.31322	6.15975	Fair
<i>Aglaia spectabilis</i> (Miq.) S.S.Jain & Bennet	Meliaceae	22	2.65147	10.6607	Poor
<i>Ailanthus integrifolia</i> Lam.	Simaroubaceae	3	0.72318	2.23857	Fair
<i>Albizia lebbek</i> (L.) Benth.	Mimosaceae	1	0.03361	0.38852	None
<i>Alseodaphne petiolaris</i> Hook. f.	Lauraceae	3	0.04198	1.0641	Fair
<i>Alstonia scholaris</i> (L.) R. Br.	Apocynaceae	1	0.01833	0.36217	None
<i>Altingia excelsa</i> Noronha	Altingiaceae	16	0.52636	5.40771	Fair
<i>Aquilaria malaccensis</i> Lam.	Thymelaeaceae	2	0.01846	0.69297	Fair
<i>Artocarpus chaplasha</i> Roxb.	Moraceae	28	6.29636	18.1394	Good
<i>Artocarpus lakoocha</i> Roxb.	Moraceae	3	0.19543	1.32867	Fair
<i>Baccaurea ramiflora</i> Lour.	Phyllanthaceae	10	0.1217	3.51555	Good
<i>Balakata baccata</i> (Roxb.) Esser	Euphorbiaceae	2	1.17837	2.69281	Fair
<i>Barringtonia acutangula</i> (L.) Gaertn.	Lecythidaceae	2	0.24522	1.08393	Fair
<i>Canarium bengalense</i> Roxb.	Burseraceae	8	0.63933	3.35239	Good
<i>Carallia brachiata</i> (Lour.) Merr.	Rhizophoraceae	3	0.0572	1.09034	Good
<i>Castanopsis indica</i> (Roxb. ex Lindl.) A.DC.	Fagaceae	20	1.72433	8.59823	Fair
<i>Castanopsis tribuloides</i> (Sm.) A.DC.	Fagaceae	8	2.38789	6.76163	Fair
<i>Chukrasia tabularis</i> A.Juss.	Meliaceae	2	0.04891	0.74547	Fair
<i>Cinnamomum glaucescens</i> (Nees) Hand.-Mazz.	Lauraceae	14	1.54341	6.50011	Poor
<i>Cyathea gigantea</i> (Wall. ex Hook.) Holtt.	Cyatheaceae	1	0.00959	0.3471	None
<i>Dillenia indica</i> L.	Dilleniaceae	4	1.70451	4.2611	Poor
<i>Diospyros variegata</i> Kurz	Ebenaceae	2	0.02531	0.70478	Good
<i>Dipterocarpus retusus</i> Bl.	Dipterocarpaceae	24	3.69504	13.91	Fair
<i>Drimycarpus racemosus</i> (Roxb.) Hook.f. ex Marchand.	Anacardiaceae	2	0.08046	0.79987	Good
<i>Duabanga grandiflora</i> (DC.) Walp.	Lythraceae	2	0.0368	0.7246	None
<i>Dysoxylum gotadhora</i> (Buch.-Ham.) Mabb.	Meliaceae	23	0.86818	7.71934	Good
<i>Elaeocarpus serratus</i> L.	Elaeocarpaceae	4	0.07487	1.45138	Fair
<i>Endospermum diadenum</i> (Miq.) Airy Shaw	Euphorbiaceae	4	0.10866	1.50962	Good
<i>Eurya acuminata</i> DC.	Pentaphylacaceae	2	0.02101	0.69737	Fair
<i>Evodia meliaefolia</i> Benth.	Rutaceae	2	0.02522	0.70463	None
<i>Ficus benghalensis</i> L.	Moraceae	2	1.12092	2.59377	Fair
<i>Ficus benjamina</i> L.	Moraceae	2	0.31396	1.20246	Fair
<i>Ficus fistulosa</i> Reinw. ex Bl.	Moraceae	19	0.29231	5.60142	Good
<i>Ficus lamponga</i> Miq.	Moraceae	8	0.55708	3.40782	Fair
<i>Ficus racemosa</i> L.	Moraceae	3	0.04018	0.86374	Poor
<i>Ficus religiosa</i> L.	Moraceae	1	0.43934	1.08805	Fair
<i>Garcinia morella</i> (Gaertn.) Desr.	Clusiaceae	3	0.09923	1.1628	Good
<i>Garcinia pedunculata</i> Roxb. ex Buch.-Ham.	Clusiaceae	3	0.0434	1.06654	Good

<i>Gmelina arborea</i> Roxb.	Lamiaceae	4	0.05154	1.41115	Fair
<i>Hydnocarpus kurzii</i> (King) Warb.	Achariaceae	4	0.11162	1.51474	Poor
<i>Ilex godajam</i> Coleb. ex Hook.f.	Aquifoliaceae	6	0.09145	2.1411	Good
<i>Khasiaclunea oligocephala</i> (Havil.) Ridsdale	Rubiaceae	11	0.97748	5.3216	Good
<i>Kydia calycina</i> Roxb.	Malvaceae	6	0.11537	2.18235	Fair
<i>Lagerstroemia speciosa</i> (L.) Pers.	Lythraceae	39	3.69519	16.3047	Fair
<i>Litsea monopetala</i> (Roxb.) Pres.	Lauraceae	12	0.21	3.53998	Good
<i>Macaranga denticulata</i> (Blume) Müll.Arg.	Euphorbiaceae	2	0.09987	0.83334	None
<i>Magnolia champaca</i> (L.) Baill. ex Pierre	Magnoliaceae	3	0.29418	1.49892	Fair
<i>Magnolia griffithii</i> Hook.f. & Th.	Magnoliaceae	3	0.53187	1.90874	Fair
<i>Magnolia hodgsonii</i> (Hook.f. & Th.) H.Keng	Magnoliaceae	12	1.39598	6.37373	Poor
<i>Magnolia hookeri</i> (Cubitt & Smith) Raju & Nayar	Magnoliaceae	29	5.13547	16.074	Fair
<i>Magnolia oblonga</i> (Wall. ex Hook.f. & Thomson) Figlar	Magnoliaceae	2	0.06037	0.568	Fair
<i>Mallotus nudiflorus</i> (L.) Kulju & Welzen	Euphorbiaceae	2	0.07335	0.78761	Fair
<i>Mangifera sylvatica</i> Roxb.	Anacardiaceae	4	0.43083	1.86786	Good
<i>Mesua ferrea</i> L.	Calophyllaceae	21	3.26135	12.3678	Good
<i>Morinda angustifolia</i> Roxb.	Rubiaceae	2	0.04877	0.74522	None
<i>Neolamarckia cadamba</i> (Roxb.) Bosser	Rubiaceae	4	1.00231	3.0504	None
<i>Olea dioica</i> Roxb.	Oleaceae	6	0.2247	2.37084	Good
<i>Palaquium obovatum</i> (Griff.) Engl.	Sapotaceae	2	0.53042	1.57566	None
<i>Premna bengalensis</i> Cl.	Lamiaceae	2	0.03218	0.71663	Fair
<i>Pterospermum acerifolium</i> (L.) Willd.	Malvaceae	2	0.02715	0.70796	None
<i>Quercus gemelliflora</i> Bl.	Fagaceae	2	0.01739	0.69113	None
<i>Saurauia roxburghii</i> Wall.	Saurauiaceae	26	0.33236	8.77342	Fair
<i>Spondias mombin</i> L.	Anacardiaceae	4	1.21775	3.42185	Fair
<i>Spondias pinnata</i> (L.f.) Kurz.	Anacardiaceae	2	0.03171	0.71582	Fair
<i>Sterculia villosa</i> Roxb.	Malvaceae	2	0.03853	0.72758	None
<i>Stereospermum chelonoides</i> (L.f.) DC.	Bignoniaceae	3	0.03336	1.04924	Good
<i>Symplocos ferruginea</i> Roxb.	Symplocaceae	4	0.00975	1.14185	Fair
<i>Syzygium kurzii</i> (Duthie) Balakr.	Myrtaceae	10	0.13052	3.53075	Good
<i>Terminalia catappa</i> L.	Combretaceae	2	0.16568	0.9468	Poor
<i>Terminalia chebula</i> Retz.	Combretaceae	3	0.03446	0.8539	Good
<i>Terminalia myriocarpa</i> Van Heurck & Müll. Arg.	Combretaceae	4	0.74261	2.60265	None
<i>Tetrameles nudiflora</i> R. Br.	Tetramelaceae	2	1.12475	2.60037	Fair
<i>Vatica lanceaefolia</i> Bl.	Dipterocarpaceae	227	6.52549	55.5215	Good
<i>Vernonia arborea</i> Buch.-Ham.	Asteraceae	3	0.08561	0.94208	None
<i>Walsura robusta</i> Roxb.	Meliaceae	2	0.78779	2.0194	None

## DISCUSSIONS

### Tree diversity

The present semi-evergreen forest patch harbours rich tree diversity. The tree species richness (75 species) recorded in this study site is higher than tropical semi evergreen forest of Manipur (Devi & Yadava 2006) and Mizoram (Lalfakawma *et al.* 2009) which recorded 17 and 32 tree species, respectively. The value is comparable to semi-evergreen and evergreen forest of Little Andaman Island, India with 83 and 84 tree species, respectively (Rasingam & Parathasarathy 2009). According to Whitmore (1984), in tropical rain forests the tree species ranges from 20 to a maximum of 223 ha<sup>-1</sup>. Tree density (750 ha<sup>-1</sup>) and basal area (58.0 m<sup>2</sup>.ha<sup>-1</sup>) recorded in tropical semi-evergreen forest of HGWLS are found to be similar (685–820 tree ha<sup>-1</sup> and 18.9 to 19.58 m<sup>2</sup>.ha<sup>-1</sup> respectively) with tropical semi evergreen forest of Manipur (Devi & Yadava 2006), tropical wet evergreen forest Namdapha National Park, northeast India (34 to 610 individuals ha<sup>-1</sup> and 7.81 to 98.58 m<sup>2</sup>.ha<sup>-1</sup>, Nath *et al.* 2005) and evergreen forest of Kalakad National Park of Western Ghats (575 to 855 individuals ha<sup>-1</sup> and 61.7 to

94.6 m<sup>2</sup>.ha<sup>-1</sup>, Parthasarathy 1999). Tree density in tropical forests varies from 245 to 859 for trees of  $\geq 30$  cm gbh (Richards 1952, Ashton 1964, Campbell *et al.* 1992) and the recorded values of the present study lies within this range. Variation in density and basal area of different forest stand may be attributed by altitudinal variation, species composition, age structure, successional stage of the forest and degree of disturbance (Swamy *et al.* 2000). The Shannon-Weiner diversity index normally varies from 1.5 to 3.5 and rarely exceeds 4.5 (Kent & Coker 1992) and is generally higher in tropical forest. The Shannon-Wiener index (3.55) recorded in the present study site is higher than tropical semi evergreen forest of Mizoram (Lalfakawma *et al.* 2009). The diversity index (H') for Indian forests ranged from 0.83 to 4.1 (Singh *et al.* 1984, Parthasarathy *et al.* 1992, Visalakshi 1995) and the value of diversity index of the present study, therefore, lies within the range and it reflects high tree diversity in the study site. Simpson's index values of different Indian tropical forests ranged from 0.03 to 0.92 (Bhuyan *et al.* 2003, Nath *et al.* 2005, Devi & Yadava 2006, Deb & Sundriyal 2011, Kushwaha & Nandy 2012) and the average value is 0.06 as reported by Knight (1975). The concentration of dominance of the study site (0.05) corresponds well with the reported range for tropical forest. Evenness index (0.82) was comparable with a report from tropical wet evergreen forest of Arunachal Pradesh (Nath *et al.* 2005) and tropical evergreen region of Meghalaya (Tynsong & Tiwari 2010). The higher evenness index value reveals more consistency in species distribution. IVI value of any species indicates the dominance of species in a mixed population and it gives a total picture of the social structure of species in a community and can be used to form an association of dominant species (Parthasarathy & Karthikeyan 1997). In the present study, it was found that *Vatica lanceaefolia* Bl., a critically endangered species, records highest IVI value emerging as the dominant tree species which was followed by *Artocarpus chaplasha*, *Lagerstroemia speciosa*, *Magnolia hookeri* and *Dipterocarpus retusus*. The observation shows that HGWLS harbours rich tree diversity providing habitat and food resources to large number of fauna. High species richness means greater diversity and which leads to a higher community stability (MacArthur 1955). However, the anthropogenic activities prevailing in the sanctuary like grazing by cattle and firewood collection by the local people to meet their energy requirements imposed threat to the survival and population structure of the species. So, if the present trend of anthropogenic pressure extended, the growth, survival and reproductive potential of the tree species will jeopardise in near future. Therefore, a proper strategy for the conservation and management of the study site is required to formulate, considering a sustainable harvest and utilization of forest resources by the local dwellers, for countering the same.

#### *Population structure and regeneration status*

The size class distribution of tree has often been used to represent the population structure of forests (Saxena & Singh 1984, Khan *et al.* 1987). Girth class frequency showed reverse J-shaped population curve in our present study which is similar to those reported from forest of North east India (Upadhaya *et al.* 2004, Mishra *et al.* 2005, Tynsong & Tiwari 2011), Eastern Ghats (Kadavul & Parthasarathy 1999, Sahu *et al.* 2012), Andaman Island (Rajkumar & Parathasarathy 2008, Rasingam & Parathasarathy 2009). The reverse J-shaped population curve of trees suggests an evolving or expanding population, climax or stable type of population in forest ecosystem, indicating that the forest harbours a growing and healthy population (Parthasarathy & Karthikeyan 1997, Mishra *et al.* 2005, Sahu *et al.* 2012). Micro-environmental factors which vary with seasonal changes have an effect on different growth stages of trees i.e. seedling, sapling, coppice and young trees that also helps to maintain the population structure (Khumbongmayum *et al.* 2006). The presence of established seedlings of dominant species like *Vatica lanceaefolia*, *Artocarpus chaplasha*, *Mesua ferrea*, etc. is an indicative of excellent recruitment of these species which also reflect that the prevailing environmental conditions of the study site are favourable for their establishment stage. High stem density in lower girth class representing young stage also reveals high biotic potential of the species which may be supported by the existing environmental conditions.

Tree regeneration can be predicted by the structure of their populations (Khan *et al.* 1987). In general, regeneration of species is affected by various anthropogenic factors (Sukumar *et al.* 1994, Khan & Tripathi 1989, Barik *et al.* 1996b, Iqbal *et al.* 2012) and natural phenomena (Welden *et al.* 1991, Iqbal *et al.* 2012). The overall regeneration status of the tree species of the study site is satisfactory at community level showing 'good' regeneration status, but 17% tree species, falls under 'not regenerating' condition may affect the population size in HGWLS in future. Species under 'not regenerating' condition might have been occurred due to existing disturbance in the study site like, grazing, firewood collection, and poor biotic potential of tree species which either affect the fruiting and seed germination or successful conversion of seedling to sapling stage. Moreover,

individuals in young stages of any species are more vulnerable to any kind of environmental stress and anthropogenic disturbance. The successful regeneration of a tree species depends on its ability to produce large number of seedlings and the ability of seedlings and saplings to survive and grow (Good & Good 1972). The forest having good canopy cover might have affected the survival of seedlings under good canopy (Pokhriyal *et al.* 2010) probably by reducing the penetration of sunlight reaching down to the forest floor. The ‘poor’, ‘none’ and ‘new’ regenerating categories include many important and useful tree species namely *Cinnamomum glaucescens*, *Dillenia indica*, *Ficus racemosa*, *Duabanga grandiflora*, *Neolamarckia cadamba*, *Syzygium cumini*, *Terminalia bellirica*, *Toona ciliata*, etc. which have certain economic values (timber and non timber forest products) and act as a source of food for the seven primate species residing in the sanctuary. 13 species contributing 15% of tree species were ‘new arrivals’/ newly colonize to the study site, representing only in sapling or seedling stage. These species may have reached or colonized to the study site by dispersal of seeds through drooping of birds and animals and getting favourable microsite to germinate and establish. Another possible reason may be that the adult individuals were very poor and have been felled by locals but seed remain as seed bank which germinate during favourable season. On the other hand, regeneration of a species is affected by various factors such as light, canopy density, soil moisture, nutrients and anthropogenic pressure (Iqbal *et al.* 2012). Small openings in the forest canopy allow higher light availability in the forest floor which favours the seedling recruitment process of certain light demanding species (Webb & Sah 2003).

## CONCLUSIONS

Precise assessment and understanding of the dynamics of plant resources is important for their sustainable management, utilization and biodiversity conservation. Quantitative analysis of tree species diversity of HGWLS will be useful in forest management and conservation as the location of the sanctuary, amidst tea gardens and villages, has made the flora more vulnerable with respect to human interference. This study provides a critical analysis of tree species richness in the study site. A reverse J-shaped population curve indicates high tree species richness and density in lower girth class which gradually decrease with increase in girth class population size. The overall population structure of tree species in the study site reveals that contribution of seedlings to the total population was highest followed by saplings and adult trees. It shows regeneration of tree species in the forest is ‘good’ and the future communities may be sustained unless there is any major environmental stress or interference exerted by human activities. However, considering the increasing anthropogenic pressure, there may be spatial and temporal threat to the seedling establishment and growth of tree species in the study site. The growth, survival and reproductive potential of the tree species will be at risk in near future if the present trend of anthropogenic continues. Thus, a systematic management plan is required for the conservation of vegetation and sustainable use of available resource. Quantitative analysis of diversity, population structure and regeneration status of tree species recorded from the present study may provide baseline information for formulating conservation and management strategies of the present forest.

## ACKNOWLEDGEMENTS

Authors acknowledge their sincere thanks to the P.C.C.F (Wildlife) and D.F.O, Jorhat Division (Assam Forest Department), for permission to carry out the work in Hollongapar Gibbon Wildlife Sanctuary. The help given by Deben Borah and other staff of Meleng Beat office during the field work is gratefully acknowledged. We are also thankful to the staff of BSI, Arunachal Pradesh Circle, Itanagar, Jintu Sarma and Monoranjan Nath for immense help and support.

## REFERENCES

- Ashton PS (1964) A quantitative phytosociological technique applied to tropical mixed rain forest vegetation. *Malaysian Forester* 27: 304–317.
- Barik SK, Tripathi RS, Pandey HN & Rao P (1996a) Tree regeneration in a subtropical humid forest: Effect of cultural disturbance on seed production, dispersal and germination. *Journal of Applied Ecology* 33: 1551–1560.
- Barik SK, Rao P, Tripathi RS & Pandey HN (1996b) Dynamics of tree seedling population in a humid subtropical forest of northeast India as related to disturbances. *Canadian Journal of Forest Research* 26: 584–589.

- Bhuyan P, Khan ML & Tripathi RS (2003) Tree diversity and population structure in undisturbed and human-impacted stands of tropical wet evergreen forest in Arunachal Pradesh, Eastern Himalayas, India. *Biodiversity and Conservation* 12: 1753–1773.
- Boerner RE & Brinkman JA (1996) Ten years of tree seedling establishment and mortality in an Ohio deciduous forest complex. *Bulletin of the Torrey Botanical Club* 123: 309–317.
- Campbell DG, Stone JL & Rosas JR A (1992) A comparison of the phytosociology and dynamics of three floodplain (varzea) forest of known ages, Rio Jurua, western Brazilian Amazon. *Botanical Journal of the Linnean Society* 108: 213–37.
- Borah N & Garkoti SC (2011) Tree species composition, diversity, and regeneration patterns in disturbed and undisturbed forests of Barak Valley, South Assam, India. *International Journal of Ecology and Environmental Sciences* 37: 131–141.
- Ceccon E, Huante P & Rincón E (2006) Abiotic factors influencing tropical dry forests regeneration. *Brazilian Archives of Biology and Technology* 49: 305–312.
- Champion HG & Seth SK (1968) *A revised survey of the forest types of India*. Govt. of India Press, New Delhi.
- Chetry D, Chetry R & Bhattacharjee PC (2007) *Hoolock: The Ape of India*. Gibbon Conservation Centre, Assam, India.
- Cottam G & Curtis JT (1956) The use of distance measurement in phytosociological sampling. *Ecology* 37: 451–460.
- Deb P & Sundriyal RC (2011) Vegetation dynamics of an old-growth lowland tropical rainforest in North-east India: species composition and stand heterogeneity. *International Journal of Biodiversity and Conservation* 3: 405–430.
- Devi LS & Yadava PS (2006) Floristic diversity assessment and vegetation analysis of tropical semievergreen forest of Manipur, north east India. *Tropical Ecology* 47: 89–98.
- Dutta G & Devi A (2013) Plant diversity, population structure, and regeneration status in disturbed tropical forests in Assam, northeast India. *Journal of Forestry Research* 24: 715 –72.
- Ganesan R & Davidar P (2003) Effect of logging on the structure and regeneration of important fruit bearing trees in a wet evergreen forest, southern Western Ghats, India. *Journal of Tropical Forest Science* 15: 12–25.
- Getachew T, Demel T, Masresha F & Erwin B (2010) Regeneration of seven indigenous tree species in a dry Afromontane forest, Southern Ethiopia. *Flora* 205: 135–143.
- Good NF & Good RE (1972) Population dynamics of tree seedlings and saplings in mature eastern hardwood forests. *Bulletin of the Torrey Botanical Club* 99: 172–178.
- Guariguata MR (1990) Landslide disturbance and forest regeneration in the upper Luquillo mountains of Puerto Rico. *Journal of Ecology* 78: 814–832.
- Guarino ESG & Scariot AO (2012) Tree seedling survival and growth in logged and undisturbed seasonal deciduous forest fragments in central Brazil. *Journal of Forest Research* 17: 193–201.
- Huston MA (1994) *Biological Diversity*. Cambridge University Press, Cambridge.
- IUCN (2014) The IUCN red list of threatened species, version 2014.1. IUCN Red List Unit, Cambridge U.K. Available from: <http://www.iucnredlist.org/> (accessed: 28 June, 2014).
- Iqbal K, Pala NA, Bhat JA & Negi AK (2012) Regeneration status of trees around Khoh river in Garhwal Himalaya. *Indian Journal of Forestry* 35: 471–476.
- Kadavul K & Parthasarathy N (1999) Plant biodiversity and conservation of tropical semi-evergreen forests in the Shervarayan hills of Eastern Ghats, India. *Biodiversity and Conservation* 8: 421–439.
- Kadavul K & Parthasarathy N (2001) Population analysis of *Alphonsea sclerocarpa* Thw. (Annonaceae) in the Kalrayan hills of Eastern Ghats, India. *International Journal of Ecology and Environmental Science* 27: 51–54.
- Kanjilal UN & Bor NL (2005) *Flora of Assam*. Omsons Publications, New Delhi.
- Kent M & Coker P (1992) *Vegetation description and analysis: A practical approach*. Belhaven Press, London.
- Khan ML & Tripathi RS (1989) Effects of stump diameter, stump height and sprout density on the sprout growth of four tree species in burnt and unburnt forest plots. *Acta Oecologia* 10: 303–316.
- Khan ML, Rai JPN & Tripathi RS (1987) Population structure of some tree species in disturbed and protected sub-tropical forests of north-east India. *Acta Oecologica-Oecologia Applicata* 8: 247–255.
- Khumbongmayum AD, Khan ML & Tripathi RS (2005) Survival and growth of seedlings of a few tree species in the four sacred groves of Manipur, Northeast India. *Current Science* 88: 1781–1788.

- Khumbongmayum AD, Khan ML & Tripathi RS (2006) Biodiversity conservation in sacred groves of Manipur, northeast India: population structure and regeneration status of woody species. *Biodiversity and Conservation* 15: 2439–2456.
- Knight DH (1975) A phytosociological analysis of species rich tropical forest on Barro-Colorado Island: Panama. *Ecological Monograph* 45: 259–289.
- Kushwaha SPS & Nandy S (2012) Species diversity and community structure in sal (*Shorea robusta*) forests of two different rainfall regimes in West Bengal, India. *Biodiversity and Conservation* 21: 1215–1228.
- Lalfakawma, Sahoo UK, Roy S, Vanlalhratpuia K & Vanlalhluna PC (2009) Community composition and tree population structure in undisturbed and disturbed tropical semi-evergreen forest stands of north-east India. *Applied Ecology and Environmental Research* 7: 303–318.
- Liang SY & Seagle SW (2002) Browsing and microhabitat effects on riparian forest woody seedling demography. *Ecology* 83: 212–227.
- MacArthur RH (1955) Fluctuation of animal population, and a measure of community stability. *Ecology* 36: 533–536.
- Mani S & Parthasarathy N (2006) Tree diversity and stand structure in inland and coastal tropical dry evergreen forests of peninsular India. *Current Science* 90: 1238–1246.
- Mishra BP, Tripathi OP & Laloo RC (2005) Community characteristics of a climax subtropical humid forest of Meghalaya and population structure of ten important tree species. *Tropical Ecology* 46: 241–251.
- Nandy S & Das AK (2013) Comparing tree diversity and population structure between a traditional agroforestry system and natural forests of Barak valley, Northeast India. *International Journal of Biodiversity Science, Ecosystem Services & Management* 9: 104–113.
- Nath PC, Arunachalam A, Khan ML, Arunachalam K & Barbhuiya AR (2005) Vegetation analysis and tree population structure of tropical wet evergreen forests in and around Namdapha National Park, Northeast India. *Biodiversity and Conservation* 14: 2109–2136.
- Negi CS & Nautiyal S (2005) Phytosociological studies of a traditional reserve forest- Thal Ke Dhar, Pithoragarh, Central Himalayas (India). *Indian Forester* 131: 519–534.
- Palit D, Pal S & Chanda S (2012) Diversity and richness of plants in Darjeeling Himalaya with an eye on Gaddikhana forest beat, Senchal east zone forest range, Darjeeling. *Indian Journal of Forestry* 35: 39–44.
- Parthasarathy N & Karthikeyan R (1997) Plant biodiversity inventory and conservation of two tropical dry evergreen forests on the Coromandel Coast, south India. *Biodiversity and Conservation* 6: 1063–1083.
- Parthasarathy N (1999) Tree diversity and distribution in undisturbed and human-impacted sites of tropical wet evergreen forest in Southern Western Ghats, India. *Biodiversity and Conservation* 8: 1365–1381.
- Parthasarathy N, Kinbal V & Kumar LP (1992) *Plant species diversity and human impact in tropical wet evergreen forests of southern Western Ghats*. Indo-French Workshop on tropical forest ecosystem: Natural Functioning and Anthropogenic Impact. French Institute, Pondichery.
- Patiri B & Borah A (2007) *Wild edible plants of Assam*. Guwahati. Director Forest Communication, Forest Department, Assam.
- Pielou EC (1966) Shannon's formula as a measure of specific diversity: its use and misuse. *American Naturalist* 100: 463–465.
- Pokhriyal P, Uniyal P, Chauhan DS & Todaria NP (2010) Regeneration status of tree species in forest of Phakot and Pathri Rao watersheds in Garhwal Himalaya. *Current Science* 98: 171–174.
- Rahman MH, Khan ASA, Roy B & Fardusi MJ (2011) Assessment of natural regeneration status and diversity of tree species in the biodiversity conservation areas of Northeastern Bangladesh. *Journal of Forestry Research* 22: 551–559.
- Rajkumar M & Parthasarathy N (2008) Tree diversity and structure of Andaman Giant Evergreen Forests, India. *Tropical Ecology* 53: 356–368.
- Ram J, Kumar A & Bhatt J (2004) Plant diversity in six forest types of Uttaranchal, Central Himalaya, India. *Current Science* 86: 975–978.
- Rannie WF (1986) Summer air temperature and number of vascular species in arctic Canada. *Arctic* 39: 133–137.
- Rasingam L & Parathasarathy N (2009) Tree species diversity and population structure across major forest formations and disturbance categories in Little Andaman Island, India. *Tropical Ecology* 50: 89–102.
- Richards PW (1952) *The tropical rain forest*. Cambridge University Press, London.

- Richards PW (1996) *The tropical rain forest: an ecological study*. - Second edition. Cambridge University Press, London.
- Sahu SC, Dhal NK & Mohanty RC (2012) Tree species diversity, distribution and population structure in a tropical dry deciduous forest of Malyagiri hill ranges, Eastern Ghats, India. *Tropical Ecology* 52: 163–168.
- Saikia P & Khan ML (2013) Population structure and regeneration status of *Aquilaria malaccensis* Lam. in homegardens of Upper Assam, northeast India. *Tropical Ecology* 54: 1-13.
- Saxena AK & Singh JS (1984) Tree population structure of certain Himalayan forest associations and implications concerning their future composition. *Vegetatio* 58: 61–69.
- Shankar U (2001) A case of high tree diversity in a sal (*Shorea robusta*)-dominated lowland forest of Eastern Himalaya: Floristic composition, regeneration and conservation. *Current Science* 81: 776–786.
- Shannon CE & Wiener W (1963) *The mathematical theory of communities*. University of Illinois Press, Urbana Illinois.
- Simpson EH (1949) Measurement of diversity. *Nature* 163: 688.
- Singh JS, Rawat YS & Chaturvedi OP (1984) Replacement of Oak forest with pine in the Himalaya affect the nitrogen cycle. *Nature* 311: 54–56.
- Sukumar R, Suresh HS, Dattaraja HS & Joshi NV (1994) In: Dallmeier F & Comiskey JA (eds) *Forest Biodiversity Research-Monitoring and Modeling, Conceptual Background to Old World Case Studies*. Parthenon publishing.
- Swamy PS, Sundarapandian SM, Chandrasekar P & Chandrasekaran S (2000) Plant species diversity and tree population structure of a humid tropical forest in Tamil Nadu, India. *Biodiversity and conservation* 9: 1643–1669.
- Tynsong H & Tiwari BK (2011) Diversity and population characteristics of woody species in natural forests and arecanut agroforests of south Meghalaya, northeast India. *Tropical Ecology* 52: 243–252.
- Tynsong H & Tiwari BK (2010) Diversity of plant species in arecanut agroforest of south Meghalaya, north-east India. *Journal Forestry Research* 21: 281–286.
- Upadhaya K, Pandey HN, Law PS & Tripathi RS (2004) Diversity and population characteristics of woody species in subtropical humid forest exposed to cultural disturbances in Meghalaya, Northeast India. *Tropical Ecology* 45: 303–314.
- Verma RK, Shadangi DK & Totey NG (1999) Species diversity under plantation raised on a degraded land. *The Malaysian Forester* 62: 95–106.
- Visalakshi N (1995) Vegetation analysis of two tropical dry evergreen forest in southern India. *Tropical Ecology* 36: 117–127.
- Ward JS, Worthley TE, Smallidge PJ & Bennett KP (2006) *Northeast forest regeneration handbook: A guide for forest owners, harvesting practitioners, and public officials*. USDA Forest Service, Newton Square, PA.
- Webb EL & Sah RN (2003) Structure and diversity of natural and managed Sal (*Shorea robusta* Gaertn. f.) forest in the Terai of Nepal. *Forest Ecology and Management* 176: 337–353.
- Welden CW, Hewett SW, Hubbell SP & Foster RB (1991) Sapling survival, growth and recruitment: relationship to canopy height in a neotropical forest. *Ecology* 72: 35–50.
- Whitmore TC (1984) *Tropical rain forest of the Far East*.- Second edition. Oxford University Press, Oxford, UK.
- Wyatt-Smith J (1987) Problems and prospects for natural management of tropical moist forests. In: Mergen F & Vincent JR (eds) *Natural Management of Tropical Moist Forests. Silvicultural and Management Prospects of Sustained Utilization* Yale University, New Haven, Connecticut, USA.